

## Rb–Sr ISOTOPIC SYSTEMATICS OF ALKALI-RICH FRAGMENTS IN YAMATO-74442 AND BHOLA.

T. Yokoyama<sup>1</sup>, K. Misawa<sup>1,2</sup>, O. Okano<sup>3</sup>, C.-Y. Shih<sup>4</sup>, L. E. Nyquist<sup>5</sup>, J. I. Simon<sup>5</sup>, M. J. Tappa<sup>4</sup>, S. Yoneda<sup>7</sup>. <sup>1</sup>SOKENDAI, Tachikawa, Tokyo 190–8518, Japan. E-mail: [yokoyama.tatsunori@nipr.ac.jp](mailto:yokoyama.tatsunori@nipr.ac.jp). <sup>2</sup>Natl Inst. Polar Res., <sup>3</sup>Okayama Univ., <sup>4</sup>ESCG/Jacobs., <sup>5</sup>NASA-JSC, <sup>7</sup>Natl Museum Natural and Sci.

**Introduction:** Alkali-rich igneous fragments were identified in the brecciated LL-chondrites, Krähenberg (LL5) [1], Bhola (LL3–6) [2], and Yamato (Y)-74442 (LL4) [3–5], and show characteristic fractionation patterns of alkaline elements [6]. The K–Rb–Cs-rich fragments in Krähenberg, Bhola, and Y-74442 are very similar in mineralogy and petrography (olivine + pyroxene + glass), suggesting that they could have come from related precursor materials [5]. We have undertaken Rb–Sr isotopic studies on alkali-rich fragments in Bhola and Y-74442 to precisely determine their crystallization ages and isotopic signatures of their precursor material(s).

**Results and Discussion:** Rubidium in alkali-rich fragments from Y-74442 is highly enriched: 20–180 times those of ordinary chondrites (OC). Alkali-rich fragments from Y-74442 yield a Rb–Sr age of  $4.429 \pm 0.054$  Ga ( $n = 9$ , error =  $2\sigma$ ) with an initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio  $I_{\text{Sr}} = 0.7144 \pm 0.0094$  ( $2\sigma$ ), indicating a young crystallization age with a high  $I_{\text{Sr}}$  compared with those of LL-chondrite whole-rocks ( $T_{\text{LL}} = 4.542 \pm 0.015$  Ga,  $I_{\text{SrLL}} = 0.69901 \pm 0.00011$ ; recalculated with  $\lambda(^{87}\text{Rb}) = 1.402 \times 10^{-11} \text{a}^{-1}$ ) [7].

Assuming that the precursor material enriched in Rb was formed at  $T_0 = 4.568$  Ga with an  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio = 0.69888 (i.e., identical to the Allende initial  $^{87}\text{Sr}/^{86}\text{Sr}$ ,  $I_{\text{SrALL}}$  [8]) and the fragments crystallized from the melt at  $T = 4.429$  Ga, a time-averaged Rb/Sr ratio is calculated to be  $\sim 2.5$ , which is much higher than the Rb/Sr ratios of the solar nebula or CI-chondrites ( $\text{Rb}/\text{Sr}_{\text{SOLAR}} = 0.65$  [8];  $\text{Rb}/\text{Sr}_{\text{CI}} = 0.32$  [9]). Any solid/liquid fractionation processes could be ruled out for K, Rb, and (Cs) enrichments since these alkali-rich fragments show flat REE patterns [6]. We suggest that a condensate from the nebular gas was a carrier of alkaline elements. It was formed from the residual nebular gas after removal of a refractory lithophile component (i.e. Sr). The temperature at which the moderately volatile component (i.e. enriched in K, Rb, and Cs) was removed from the gas was relatively high ( $\geq 970$  K, the 50% condensation temperature of Na [10]). This component was isolated from other lithophiles and might have been isolated from the nebular gas without any interaction with high/low temperature components. At 4.43 Ga the condensate was incorporated into ferromagnesian components (i.e., olivine+pyroxene) possibly during an impact melting event. They suffered further fragmentations due to impacts and then were incorporated into the LL-chondritic breccia.

**References:** [1] Kempe W. and Müller O. (1969) *Meteorite Res.*, pp. 418. [2] Noonan A. F. *et al.* (1978) *Geol. Survey Open File Report*, **78–701**, 311. [3] Yanai K. *et al.* (1978) *Mem. Natl. Inst. Polar Res. Spec. Issue*, **8**, 110. [4] Ikeda Y. and Takeda H. (1979) *Mem. Natl. Inst. Polar Res. Spec. Issue*, **15**, 123. [5] Yokoyama T. *et al.* (2011) *LPS XLII*, Abstract #1941. [6] Wlotzka F. *et al.* (1983) *Geochim. Cosmochim. Acta*, **47**, 743. [7] Minster and Allègre (1981) *Erath and Planet. Sci. Lett.*, **56**, 89. [8] Gray C. M. *et al.* (1973) *Icarus*, **20**, 213. [9] Palme H. (2001) *Phil. Trans. R. Soc. Lond. A* **2001**, 359, 2061. [10] Wasson J. T. (1985) *Meteorite*, pp 267. Freeman, NY.